Patient Understanding of Diabetes Self-Management: Participatory Decision-Making in Diabetes Care

Charlene C. Quinn, R.N., Ph.D.,¹ Renee Royak-Schaler, Ph.D.,² Dan Lender, M.D.,³ Nanette Steinle, M.D.,³ Shahinaz Gadalla, M.D., Ph.D.,² and Min Zhan, Ph.D.²

Abstract

Objective:

Our aim was to determine whether patient participation in decision-making about diabetes care is associated with understanding of diabetes self-management and subsequent self-care practices. We also identified issues that would impact messaging for use in mobile diabetes communication.

Research Design and Methods:

A cross-sectional observational study was conducted with type 2 diabetes patients (n = 81) receiving their care at the University of Maryland Joslin Diabetes Center. A convenience sample of patients were eligible to participate if they were aged 25–85 years, had type 2 diabetes, spoke English, and visited their physician diabetes manager within the past 6 months. In-person patient interviews were conducted at the time of clinic visits to assess patient understanding of diabetes management, self-care practices, and perceptions of participation in decision-making about diabetes care.

Results:

African Americans reported fewer opportunities to participate in decision-making than Caucasians, after controlling for education [mean difference (MD) = -2.4, p = .02]. This association became insignificant after controlling for patient–physician race concordance (MD = -1.5, p = .21). Patient understanding of self-care was predicted by having greater than high school education (MD = 3.6, p = .001) and having physicians who involved them in decision-making about their care. For each unit increase in understanding of diabetes self-care, the mean patient self-care practice score increased by 0.16 (p = .003), after adjustment for patient race and education.

Conclusions:

Patient participation in decision-making is associated with better understanding of care. Participation in decision-making plays a key role in patient understanding of diabetes self-management and subsequent self-care practices. Patients with limited education need specific instruction in foot care, food choices, and monitoring hemoglobin A1c.

J Diabetes Sci Technol 2011;5(3):723-730

Author Affiliations: ¹Department of Epidemiology and Public Health, Division of Gerontology, University of Maryland School of Medicine, Baltimore, Maryland; ²Department of Epidemiology and Public Health, University of Maryland School of Medicine, Baltimore, Maryland; and ³Department of Medicine, Division of Endocrinology, University of Maryland School of Medicine, Baltimore, Maryland

Abbreviations: (CI) confidence interval, (HbA1c) hemoglobin A1c, (MD) mean difference, (OR) odds ratio, (PDM) participatory decision-making

Keywords: literacy, mobile, participatory, self-care

Corresponding Author: Charlene C. Quinn, R.N., Ph.D., University of Maryland School of Medicine, Department of Epidemiology and Public Health, 660 W. Redwood St., Baltimore, MD 21201; email address <u>cquinn@epi.umaryland.edu</u>